

RESEARCH ARTICLE

Open Access



DELAY on first antenatal care visit and its associated factors among pregnant women in public health facilities of Debre Markos town, North West Ethiopia

Atsede Alle Ewunetie^{1*}, Alemtsehay Mekonnen Munea², Belsity Temesgen Meselu³, Muluye Molla Simeneh¹ and Bekele Tesfaye Meteku⁴

Abstract

Background: Delay on timely initiation of antenatal care has a great impact on adverse pregnancy outcomes. However, evidences in Ethiopia revealed that majority of pregnant mothers did not start their first visit as recommended by WHO. The aim of this study was to assess delay and associated factors of first antenatal care visit among pregnant mothers at public health facilities of Debre Markos town, North West Ethiopia.

Methods: An institutional based cross-sectional study was conducted from February to March, 2014 in public health facilities of Debre Markos town North west Ethiopia. A total of 320 pregnant mothers who were sure of their last menstrual periods were interviewed with a structured questionnaire. Data entry was done using Epi data 3.1 and analysis was done using SPSS version 20. Descriptive statistics, binary and multivariable logistic regression analyses were employed to identify the magnitude and factors associated with delay on timely initiation of the first antenatal care visit.

Results: The proportion of respondents who made their first antenatal care visit after 16 weeks of gestation was found to be 33.4%. Mothers residing in rural settings (AOR = 2.8 [95% CI:1.54–5.44]), not attained formal education (AOR = 2.2 [95% CI:1.10–4.68]), with unintended pregnancy (AOR = 3.6 [95% CI:2.00–6.80]) and who perceived that the right initiation time of the first antenatal care visit is beyond 16 weeks of gestation (AOR = 3.9 [95% CI:1.61–9.76]) were more likely delayed on their first antenatal care visit.

Conclusion: Residence, educational status, intention of pregnancy and perception on the right time of first antenatal care visit initiation were found to be predictors of delay on timely initiation of first antenatal care visit. Therefore, the Zonal health department should strengthen awareness creation about timely initiation of first antenatal care visit and family planning to prevent unintended pregnancy in the community especially in the rural settings.

Keywords: Delay on timely initiation of antenatal care, Unintended, Pregnancy

* Correspondence: atsiale@gmail.com

¹Department of Public Health, College of Health Sciences, Debre Markos University, Debre Markos, Ethiopia

Full list of author information is available at the end of the article



Background

Many health problems of pregnant women are preventable, detectable, and treatable if they attained antenatal care (ANC) adequately [1]. The traditional approach of the ANC is a high risk approach which intended to classify pregnant women at low risk or high risk based on predetermined criteria and involved many ANC visits. This approach was hard to implement effectively since many women had at least one risk factor, and not all developed complications; at the same time, some low risk women did develop complications. It is a burden on the healthcare system. As a result, many developing countries, including Ethiopia, are adopting focused antenatal care (FANC) approach. The newly proposed, FANC recommends four ANC visits for most pregnant women. Ideally, the first visit is in the first trimester but not after 16 weeks of gestation [2, 3].

The recommendation sets out from the fact that first trimester pregnancy stage is the fastest developmental period of the fetus, in which all organs become well developed and needs special attention [4, 5]. However, too many women make their first antenatal visit with the pregnancy already compromised or at risk from smoking, inappropriate nutrition, ingestion of a variety of drugs, including pharmaceutical preparations, genitourinary tract infections, anemia and poor dental health [5].

Women present for antenatal care early in their pregnancy period allow enough time for essential and feasible interventions, prevention of complications and early identification of underlying conditions [6]. It also used to prevent, diagnose and treat sexually transmitted infections and work on the elimination of new Human Immune Deficiency Virus infections among new borne through providing integrated quality prevention of mother to child transmission [7–9].

Early attendance of ANC provides a better hemoglobin concentration through nutritional advice, prevention and early treatment of malaria and timely iron foliate supplementation [10, 11]. Beside this, it increases the opportunity of pregnant mothers to have more prenatal care visits, sufficient tests and advice during pregnancy and a skilled birth attendant [12, 13].

Mothers who start ANC after 22 weeks gestation, missed over four routine antenatal visits, who did not seek care or who concealed their pregnancy were manifest 17% of maternal deaths [14]. Suboptimal antenatal care was also found to be the major contributory factor for still birth in India [15]. Mothers who sought antenatal care before the end of the third month had infants who weighed heavier compared to the infants of mothers who sought care later [11, 16, 17].

Even though there is improvement on antenatal care coverage and the World Health Organization recommendation is initiating ANC visit in the first trimester,

the time of initiation of first ANC visit is varied throughout the world. In the Ethiopian context, it is recommended that the first ANC visit should be ideally taken place before 16 weeks of pregnancy (2). In order to improve maternal health care service utilization; all governmental health institutions of Ethiopia are providing focused antenatal care service for all pregnant mothers who come to the health institutions free of charge. Early initiation of antenatal care is promoted by health extension workers and health professionals in both urban and rural Keble's. Beside women developmental armies who are delegated in the community have their own role in community mobilization regarding to antenatal care service utilization.

According to the Ethiopia demographic, health survey, over 34% of pregnant women were attending antenatal care at least once. But, only 19% had four or more antenatal care visits during their entire pregnancy, and 11% of women made their first antenatal care visit before the fourth month of pregnancy nationwide [3]. In order to improve maternal and child health, identifying timing of first antenatal care visit among pregnant mothers and factors that affect initiation of first ANC visit is paramount. This study is designed to assess delay on first antenatal care visit among pregnant mothers and factors that affect initiation of first ANC.

Methods

Study design and data sources

An institutional based cross sectional study design was employed from February 1st through March 30, 2014 in public health facilities of Debre markos town. Debre markos town is the capital of East Gojjam Zone, which is located 300 km North West of Addis Ababa and 265 km away from regional capital Bahir Dar.

The study populations were pregnant women who visited public health facilities of Debre markos town for antenatal care and randomly selected during the study period. Those pregnant mothers who were seriously ill or not aware their menstrual period were excluded from the study. The sample size was determined using single population proportion formula with the following assumptions: the proportion of delayed ANC is 74% from previous study conducted in Debre Birhan (18), maximum acceptable marginal error of 5%, an alpha level of 0.05 and a none response rate 10% were used to obtain a sample size of 326.

All public health facilities (one referral Hospital and three health centers) in Debre markos town were included in the study. The total sample size was proportionally allocated to health facilities based on average monthly flow of pregnant mothers for ANC in each health facility. The study subjects were recruited by using systematic random sampling technique after identification of the first study

subject by simple random sampling method. Multiple enumerations due to referral were avoided using filtering questions.

The outcome variable of the study was delay on initiation of first ANC visit and the explanatory variables included Socio-demographic factors (age, religion, ethnicity, marital status, educational status, average monthly income), Obstetric factors (parity, History of obstetric complication, history of Previous ANC visit and intention of pregnancy), Enabling factors (accessibility of information about ANC and interaction with health professional) and Reinforcing factors (perception of mothers on the advantage of the ANC, timing of ANC visit and frequency of the ANC; reasons that initiate mothers to start ANC after 16 weeks, a decision made to seek ANC, intention of partner to wait for ANC service utilization).

Delay on initiation of first antenatal care visit is initiation of first ANC visit in public health facilities which have skilled health personnel after 16 weeks of gestation.

Data were collected using structured interviewer administered questionnaire adopted from literatures and contextualized to the local situations and study objectives. The data collection tool was translated into the local language (Amharic) and pretested on 5% of the actual sample size out of the study area. Four diploma nurses and one public health professional were involved as data collector and supervisor respectively after taking 1 day training.

Ethical considerations

The study was conducted after obtaining ethical clearance from Bahir Dar University, College of medicine and Health sciences, Research Ethics Committee and letter of support from Amhara regional health bureau ethical review committee. Formal Permission paper was given to woreda health office, Debremarkos Referral Hospital and responsible persons in each health institution accordingly. Verbal informed consent also obtained from the study participants after explaining the purpose of study. Participants were informed on their full right to skip any question or terminate their participation at any stage. Participants were also assured that there will be no harm or benefit of being participating in this study. All the information from the respondents was kept confidential.

Data processing and analysis

The data were cleaned, coded and entered in Lauritsen JM, Bruus M, Myatt M. Epi Data. A comprehensive tool for validated entry and documentation of data. 2003; 3 and transferred to IBM SPSS Statistics for Windows, Version 20.0. Armonk, NY: IBM Corp for analysis. Both descriptive and inferential statistics were used to summarize the data. Those variables found to be statistically significant

($p < 0.05$) in the binary logistic regression analyses were entered into the multivariable logistic regression model and statistical significance was considered at $p < 0.05$. The strength of association was assessed by odds ratio (OR) with 95% confidence interval.

Results

The response rate was 98.1%. The mean (\pm SD) age of respondents was 26.3 (\pm 5.1). Among all respondents, 311 (97.2%) were Amhara in ethnic group and majority 299 (93.4%) of mothers were orthodox Christian followers (Table 1).

Of all respondents, 107 (33.4%) made their first ANC visit after 16 weeks of gestation. The mean time was 14.5 (\pm 6.5) weeks with the range of 4 to 36 weeks. The median and the pick time was 12 weeks of gestation. Above

Table 1 Socio demographic characteristics of pregnant mothers who were attending ANC in the public health facilities of Debremarkos town, February-March 2014 ($n = 320$)

Variable		Frequency	Percent
Residence	Rural	69	21.6
	Urban	251	78.4
Age	15–19	16	5.0
	20–24	104	32.5
	25–29	121	37.8
	30–34	55	17.2
	≥ 35	24	7.5
Ethnicity	Amhara	311	97.2
	Others	9	2.8
Religion	Orthodox christian	299	93.4
	Muslim /Protestant	21	6.6
Marital status	Never married	14	4.4
	Married	293	91.6
	Divorced/widowed	13	4
Educational status	Have no formal education	98	30.6
	Primary education	51	15.9
	Secondary education	87	27.2
	Tertiary education	84	26.3
Occupation	Government employee	81	25.3
	House wife	49	15.3
	Merchant	118	36.9
	Farmer	42	13.1
	Daily labourer	17	5.3
Monthly income	Students / depende on family	13	4.1
	< 1000 ETB(<Q1)	124	38.8
	1000–2987.50ETB	116	36.2
> 2987.50ETB(>Q3)	80	25.0	

Others = Oromo/Tigrig/Agew

half (51.2%) of mothers were start their first ANC visit in the first trimester (Fig. 1).

Among the total respondents, 174 (54.4%) were multiparous. Of these, 126 (74.2%) reported that they had had experience of ANC for the preceding pregnancy. For those who had had ANC visit on preceding pregnancy, 43 (34%) visited the ANC clinic for the first time after 16 weeks of gestation in the current pregnancy. From 68 (21.2%) of mothers who had unintended pregnancy, 25 (36.8%) indicated that they had interest in interrupting their pregnancy (Table 2).

The available health facilities were health centers for 220 (68.8%) of respondents, Hospital for 60 (10.8%) of respondents and health posts for 40 (12.4%) of respondents. Of all respondents, 182 (56.9%) were advised for ANC on current pregnancy. Of these, only 38 (20.9%) had information on the time of first ANC visit. From those who had information on initiation time of ANC, 2 (5.3%) were informed to start their first visit after 16 weeks of gestation (Table 3).

A majority, 27 (85.3%) of respondents perceived that the ANC is important for both maternal and child health. Among delayed respondents 18 (16.8%) were initiated ANC by considering the time of their visit at the right time (Fig. 2).

Among respondents 264 (82.5%) were decided to seek ANC service jointly with their partners. From all, 287 (89.4%) partners supported the need of ANC for every pregnant mothers.

The bivariate analysis showed that pregnant mothers who were residing in rural areas [Crude Odds Ratio (COR) = 3.4, 95% Confidence Interval (CI): 2.01–6.07], not attained formal education [COR = 4.5, 95% CI: 2.33–8.67], farmers [COR = 7.8, 95% CI: 3.37–18.05], having less than 1000 ETB average monthly income [COR = 2.7, 95% CI: 1.45–5.37], experience




Fig. 1 Distribution of first ANC visits among pregnant mothers in public health facilities in DebreMarkos in each trimester between February 2014 and March 2014

Table 2 Obstetric history of pregnant mothers, who were attending ANC in the public health Facilities of Debre Markos town, Feburary – March, 2014 (n = 320)

Variable	Frequency	Percent	
Gravidity (n = 320)	Primigravida	146	45.6
	Multi gravida	174	54.4
Parity (n = 320)	No parity	158	49.4
	≥1 parity	162	50.6
Children born alive (n = 174)	0	17	9.8
	1–3	135	77.6
	≥4	22	12.6
Death of children (n = 157)	Yes	30	19.1
	No	127	80.9
spontaneous abortion (n = 174)	Yes	35	20.1
	No	139	79.9
stillbirth (n = 174)	Yes	18	10.3
	No	156	89.7
History of ANC (n = 174)	Yes	126	72.4
	No	48	27.6
ANC initiation time on Previous pregnancy (n = 126)	≤16 Weeks	115	91.3
	> 16 Weeks	11	8.7
Type of pregnancy N = 320	Planned	252	78.8
	Unplanned	68	21.2
Intended to intrupt unplanned Pregnancy N = 68	Yes	25	36.8
	No	43	63.2

Table 3 Availability and accessibility of health information among mothers who were attending ANC in the public health facilities of DebreMarkos town, Feburary-March 2014 (n = 320)

Variable	Frequency	Percent	
Health education about ANCon previous pregnancy (n = 126)	Yes	85	67.5
	No	41	32.5
Advised for ANC (n = 320)	Yes	182	56.9
	No	138	43.1
By whom you were advised (n = 182)	Husbands	105	57.7
	Health extension Workers	28	15.4
	Relatives	20	11
	Friends/nighbours	29	15.9
Advised on initiation time (n = 128)	Yes	38	20.9
	No	144	79.1
Advised time (n = 38)	≤16 weeks	36	94.7
	> 16 weeks	2	5.3




Fig. 2 Reasons for delay on first ANC visit among pregnant mothers who attended ANC in the public health facilities of Debre markos town between February 2014 and March 2014

one or more births [COR = 1.9, 95% CI: 1.22–3.15], not planned their pregnancy [COR = 3.9, 95% CI: 2.25–6.89], having a history of stillbirth [COR = 2.6, 95% CI: 1.01–6.90] and those who perceived that the right time of first ANC visit is after 16 weeks of gestation [COR = 3.8, 95% CI: 1.69–9.76] were more likely delayed on initiation of the first ANC visit. After adjusting for the potential cofounders; multivariable logistic regression analysis indicated that residence, educational status, intention of pregnancy and perception on the right time of first ANC visit were significantly associated with delayed first ANC visit.

Mothers who were residing in rural areas, not attending formal education, having unplanned pregnancy and perceiving the right time of ANC initiation as greater than 16 weeks of gestation were more likely delayed on first ANC visit (Adjusted Odds Ratio (AOR) = 2.8 [95% CI: 1.54–5.44]), (AOR = 2.2 [95% CI: 1.10–4.68]), (AOR = 3.6 [95% CI: 2.00–6.80]) and (AOR = 3.9 [95% CI: 1.61–9.76]) respectively (Table 4).

Discussion

For many of the essential interventions in ANC, it is crucial to have early identification of underlying conditions. The first ANC visit should be as early as possible in pregnancy [10]. However, evidence in Ethiopia indicated that delayed ANC initiation time among pregnant mothers was high [18–21]. In our study, one third 107 (33.4%) of respondents were delayed to start their first ANC visit with in the first 16 weeks of pregnancy. This finding is higher as compared to a study done in Bengazie (27%, 2007) [22]. This might be due to difference in educational status of mothers between Bengazie and ours. But this finding is comparatively lower than the findings studied in Ndola (68.6%) and Mpongwe (72%) districts of Zambia [23] and South East Tanzania (81.5%) [24], EDHS, 2011 (89%) [25], Jimma University Specialized Hospital (60.1%) [26], Kembata tembaro Zone (68.6%) [27] and Dembech district of East

Gojjam Zone, North west Ethiopia (94.2%) [28]. The possible explanation for this observed difference might be due to study population composition in which proportion of mothers residing in rural areas of current study were lower than other studies.

The finding also showed a significant association of residence and delay on timely initiation of ANC. Those mothers who were residing in rural areas were 2.8 times more likely delayed than urban mothers. Among all respondents, 56% of rural and 27% of urban residents were start ANC after 16 weeks of gestation. Ethiopian demographic and health survey, 2011, revealed that urban mothers made their first visit earlier (4.4 months) than rural mothers (5.5 months) [25]. This finding is also in agreement with the study done in Vietnam in which rural mothers were attained ANC latter and used fewer visits [29].

The possible reason might be better educational status of urban mothers than rural mothers. Because, in our study, 60% of rural mothers were not attained formal education compared with 23% of urban mothers. The other reason might be availability of alternative health care facilities and having a better chance of health information in urban areas than rural areas. However, in contrary to our result, there was no significant difference in the proportion of delayed ANC attendance between urban and rural areas of Zambia [23]. The reason for this difference might be the presence of more active mobile maternity service in rural than urban areas of Zambia.

Educational status was found to have significant association with delayed initiation of first ANC visit. Those mothers who had no formal education were delay two times more likely than those who had tertiary education. This is consistent with studies done in Gondar, Kembata tembaro zone, Tanzania and Ghana [11, 24, 27, 30] in which women who had lower education or none booked later than those with higher education. The reason might be high chance of exposure for information in case of educated mothers.

Table 4 Factors associated with delay on first ANC visit among pregnant mothers who were attending ANC in public health institutions of Debreworkos town, 2014 (n = 320)

Variable	Delayed		COR(CI)	AOR(CI)	P-Value
	Yes (%)	No (%)			
Residence					
Rural	39(12.2)	30(9.4)	3.4(2.01–6.07)	2.8(1.54–5.44) ^a	0.001
Urban	68(21.2)	183(57.2)	1.0	1.0	
Educational status					
Had no formal education	54(16.9)	44(13.8)	4.5(2.33–8.67)	2.2 (1.10–4.68) ^a	0.027
Primary education	16(5)	35(10.9)	1.6(0.76–3.68)	1.4(0.65–3.42)	
Secondary education	19(5.9)	68(21.3)	1.0(0.49–2.12)	0.8(0.40–1.87)	
Tertiary education	18(5.6)	66(20.6)	1.0	1.0	
Occupation					
Government employee	18(5.6)	63(19.7)	1.0	1.0	
Merchant	12(3.8)	37(11.6)	1.1(0.49–2.61)	0.9(0.30–2.98)	
House wife	34(10.6)	84(26.3)	1.4(0.73–2.73)	1.0(0.37–2.94)	
Farmer	29(9.1)	13(4.1)	7.8(3.37–18.05)	2.2(0.52–9.51)	
Daily labourer	9(2.8)	8(2.5)	3.9(1.32–11.67)	1.5(0.33–7.21)	
Student / depend on family	5(1.6)	8 (2.5)	2.1(0.63–7.51)	1.1(0.22–5.69)	
Monthly Income					
< 1000 ETB	51(15.9)	73(22.8)	2.7(1.45–5.37)	1.5(0.66–3.59)	
1000–2987.50 ETB	40(12.5)	76(23.8)	2.1(1.07–4.10)	1.4(0.66–3.30)	
> 2987.50 ETB	16(5)	64(20)	1.0	1.0	
Parity					
No parity	41(12.8)	117(36.6)	1.0	1.0	
≥ 1 parity	66(20.6)	96(30)	1.9(1.22–3.15)	1.4(0.86–2.56)	
Types of pregnancy					
Planned	67(20.9)	185(57.8)	1.0	1.0	0.000
Unplanned	40(12.5)	28(8.8)	3.9(2.25–6.89)	3.6(2.00–6.80) ^a	
History of still birth					
No still birth	97(30.3)	205(64.1)	1.0	1.0	
≥ 1 still birth	10 (3.1)	8(2.5)	2.6(1.01–6.90)	1.6(0.54–5.10)	
Perceived time for 1st ANC visit					
≤ 16 weeks	90(28.1)	203(63.4)	1.0	1.0	0.003
> 16 weeks	17(5.3)	10 (3.1)	3.8(1.69–8.70)	3.9(1.61–9.76) ^a	

^aStatistical significant at 5% alpha level

Eventhough parity had no association with delayed initiation of ANC in our study, 40.7% of multiparous and 25.9% of nuliparous mothers were start thier first ANC visit after 16 weeks of gestation. But, it was one of the factors for delayed initiation of ANC in different studies [13, 31, 32].

Intention of pregnancy was significantly associated with delay on initiation of first ANC visit. In this study, women with un planned pregnancy were 3.6 times more likely delayed to initiate first ANC visit than those mothers with planned pregnancy. This finding was inline with a study done in Kembata tembaro Zone [26]. Findings in different

studies also in agreement with the association of unplanned pregnancy and delayed initiation of first ANC visit [18, 32–35]. The possible reason might be mothers having intended pregnancy are much cautious and eger to know their pregnancy status and less likely delayed than those who had unintended pregnancy.

Intention of abortion in case of unintended pregnancy might also increased the chance of delayed initiation of ANC. Because, above half 14(56%) of pregnant mothers with unintended pregnancy, who looked for abortion did their first ANC visit after 16 weeks of gestation in our study.

Perception on the right time of ANC initiation was found to be significantly associated with delayed ANC visit. Those mothers who perceived that the right initiation period of first ANC visit is beyond 16 weeks of gestation, were four times delayed than their counter part.

The Possible reason might be decreased exposure for information related to ANC initiation time. Because, in this study only 38(11.8%) of mothers have got information about initiation time of ANC from other persons. Other similar studies also suggested that proper information and advice on pattern of ANC utilization is important to book early [33]; where as, not knowing the right gestational age at which to start the first antenatal care visit was the commonest reason for late ANC attendance [36]. So, appropriate perception of the initial ANC visit was a factor for an early ANC visit. This is in agreement with a study done in Gondar University Hospital [30].

Limitation of the study

The study design is not strong enough to identify determinant factors.

Conclusion

The magnitude of delay on initiation of first ANC visit was still high but it is lower than studies conducted in other areas of Africa and Ethiopia. The time of initiation was ranges from first trimester to third trimester and the mean gestational age on initiation of first ANC was around second trimester. According to this study residence, educational status, intention of pregnancy and perception on initiation time of ANC were influencing mothers on timely initiation of first ANC visit. Awareness creation towards timely initiation of first ANC visit and family planning utilization to prevent unplanned pregnancy should be strengthened by Zonal health department in the community specially for the reproductive age group and rural residents. Ministry of education should also improve women education by strengthening adult education in the community. Further study is recommended in the rural community.

Acknowledgements

First and for most thanks to the Almighty GOD who is our power. Our gratitude extends to East Gojjam Zone Plan and Economy department, East Gojjam zone health department, Debre markos town administrative health office and health facilities. We would like to thank Bahir Dar University and Debre Markos University for their financial and technical support. We are also grateful to data collectors and study participants for giving us their valuable time.

Availability of data and materials

All the data supporting the study findings are within the manuscript. Additional detailed information and raw data are available from the corresponding author on reasonable request.

Authors' contributions

AA drafted the proposal, did the analysis, wrote the results and prepared the manuscript. AM participated on editing, analysis and write up of the result. BT, MM and BT were involved on data analysis and manuscript preparation. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The study was conducted after obtaining ethical clearance from Bahir Dar University, College of medicine and Health sciences, Research Ethics Committee and letter of support from Amhara regional health bureau ethical review committee. Formal Permission paper was given to woreda health office, Debre markos Referral Hospital and responsible persons in each health institution accordingly. Verbal informed consent also obtained from the study participants after explaining the purpose of study. Participants were informed on their full right to skip any question or terminate their participation at any stage. Participants were also assured that there will be no harm or benefit of being participating in this study. All the information from the respondents was kept confidential.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Public Health, College of Health Sciences, Debre Markos University, Debre Markos, Ethiopia. ²Department of Public Health, College of Medicine and Health Science, Bahir Dar University, Bahir Dar, Ethiopia. ³Department of Midwifery College of Health Sciences, Debre markos University, Debre Markos, Ethiopia. ⁴Department of Nursing, College of Health Sciences, Debre Markos University, Debre Markos, Ethiopia.

Received: 13 November 2017 Accepted: 19 April 2018

Published online: 16 May 2018

References

- World Health Organization. Global health observatory (GHO) data. URL. Available form: http://www.who.int/gho/publications/world_health_statistics/2011/en/. 2015 Mar.
- Ethiopian Federal Ministry of health. Antenatal Care Module: Providing Focused antenatal care, FMOH. <http://www.open.edu/openlearncreate/modulecontent/view.php?id=44&printable=1>.
- Khanna J, Bergsjö P, Lashley K, Peters C, Sherratt D, Villar J. WHO antenatal care randomized trial: manual for the implementation of the new model. 2002. http://apps.who.int/iris/bitstream/10665/42513/1/WHO_RHR_01_30.pdf.
- WHO. Essential Antenatal, perinatal and post partum care training module :promoting effective preinatal care. Copenhagen: World Health Organization regional office for Europe; 2002.
- Abou-Zahr CL, Wardlaw TM, World Health Organization. Antenatal care in developing countries: promises, achievements and missed opportunities: an analysis of trends, levels and differentials, 1990–2001. 2003.
- FMOH. Accelerated Plan for Scaling up PMTCT Services in Ethiopia, Federal Ministry Of Health, Addis Ababa, Ethiopia. 2014.
- Bollini P, Quack-Lötscher K. Guidelines-based indicators to measure quality of antenatal care. *J Eval Clin Pract*. 2013;19(6):1060–6.
- World Health Organization. Global update on HIV treatment 2013: results, impact and opportunities. 2013.
- East Gojjam zone health department: Annual report of East Gojjam zone health department, Debre markos, Ethiopia. 2013. Unpublished local report.
- Lawn J, Kerber K. Opportunities for African newborns: practical data policy and programmatic support for newborn care in Africa. 2006.
- Tayie FA, Lartey A. Antenatal care and pregnancy outcome in Ghana, the importance of women's education. *African Journal of Food, Agriculture, Nutrition and Development*. 2008;8(3):291–303.
- Hueston WJ, Gilbert GE, Davis L, Sturgill V. Delayed prenatal care and the risk of low birth weight delivery in US. *J Community Health*. 2003;28(3):199–208.
- Abeje G, et al. Factors associated with institutional delivery service utilization among mothers in Bahir Dar city administration, Amhara region. *Reprod Health*. 2014;11(22):3–7.

14. Bragg R, Action M. Money and Maternity: charging vulnerable pregnant women for NHS care. *Maternity Action and Medact*. 2013;26:1-9.
15. Saxena V, Bansal S, Chaturvedi J, Kalra BP, Chandra V, Kansal S. Investigating causes and factors associated with stillbirth by verbal autopsy in Uttarakhand. *Indian J Prev Soc Med*. 2011;42:14-8.
16. Alibekova R, Huang JP, Chen YH. Adequate prenatal care reduces the risk of adverse pregnancy outcomes in women with history of infertility: a nationwide population-based study. *PLoS One*. 2013;8(12):e84237.
17. Gebremariam A. Factors predisposing to low birth weight in JimmaHospital southwestern Ethiopia. *East Afr Med J*. 2005;8(1):554.
18. Zegeye AM, Bitew BD, Koye DN. Prevalence and determinants of early antenatal care visit among pregnant women attending antenatal care in Debre Berhan health institutions, Central Ethiopia. *Afr J Reprod Health*. 2013;17(4):130-6.
19. Pell C, Meñaca A, Were F, Afrah NA, Chatio S, Manda-Taylor L, Hamel MJ, Hodgson A, Tagbor H, Kalilani L, Ouma P. Factors affecting antenatal care attendance: results from qualitative studies in Ghana, Kenya and Malawi. *PLoS One*. 2013;8(1):e53747.
20. Kebede B, Gebeyehu A, Andargie G. Use of previous maternal health services has a limited role in reattendance for skilled institutional delivery: cross-sectional survey in Northwest Ethiopia. *Int J Womens Health*. 2013;5:79.
21. Fenta M. Assessment of factors affecting utilization of maternal health care services in Ayssaita and Dubti towns, afar regional state, north East Ethiopia. Master thesis, 2005.
22. Ziyo FY, Matly FA, Mehemd GM, Dofany EM. Relation between prenatal care and pregnancy outcome at Benghazi. *Sudanese J Public Health*. 2009;4(4): 403-10.
23. Banda I, Michelo C, Hazemba A. Factors associated with late antenatal care attendance in selected rural and urban communities of the Copperbelt province of Zambia. *Med J Zambia*. 2012;39(3):29-36.
24. Gross K, Alba S, Glass TR, Schellenberg JA, Obrist B. Timing of antenatal care for adolescent and adult pregnant women in South-Eastern Tanzania. *BMC Pregnancy Childbirth*. 2012;12(1):16.
25. Demographic E. Health survey (2011) Addis Ababa. Ethiopia and Calverton: Central Statistical Agency and ICF International; 2012.
26. Ewnetu W, Assegid S, Wondafrash W, Ewnetu H. Factors associated with late antenatal care initiation in an Ethiopian clinic. Research. 2015.
27. Tekelab T, Berhanu B. Factors associated with late initiation of antenatal care among pregnant women attending antenatal Clinic at Public Health Centers in Kembata Tembaro zone, southern Ethiopia. *Sci Technol Arts Res J*. 2014;3(1):108-15.
28. Gedefaw M, Muche B, Aychiluhem M. Current status of antenatal care utilization in the context of data conflict: the case of Dembecha District, Northwest Ethiopia. *Open J Epidemiol*. 2014;4(04):208.
29. Toan TK. Antenatal and delivery care utilization in urban and rural contexts in Vietnam. A study in two health and demographic surveillance sites. Gothenburg, Sweden. 2012.
30. Belayneh T, Adefris M, Andargie G. Previous early antenatal service utilization improves timely booking: cross-sectional study at university of Gondar hospital, northwest Ethiopia. *J Pregnancy*. 2014;2014:1-7.
31. UmberJalil B, Yasmeen K, Razia N. Relation ship between maternal hemoglobin and prenatal out come, Pakistan railway hospital ,Rawalpindi. *Rawal Med J*. 2007;32:102-4.
32. Exavery A, Kanté AM, Hingora A, Mbaruku G, Pemba S, Phillips JF. How mistimed and unwanted pregnancies affect timing of antenatal care initiation in three districts in Tanzania. *BMC Pregnancy Childbirth*. 2013;13(1):35.
33. Tariku A, Melkamu Y, Kebede Z. Previous utilization of service does not improve timely booking in antenatal care: cross sectional study on timing of antenatal care booking at public health facilities in Addis Ababa. *Ethiopian Journal Health Dev*. 2010;24(3):227-233.
34. Birmeta K, Dibaba Y, Woldeyohannes D. Determinants of maternal health care utilization in HOLETA town, Central Ethiopia. *BMC Health Serv Res*. 2013;13(1):256.
35. Dibaba Y, Fantahun M, Hindin MJ. The effects of pregnancy intention on the use of antenatal care services: systematic review and meta-analysis. *Reprod Health*. 2013;10(1):50.
36. Kisuule I, Kaye DK, Najjuka F, Ssematimba SK, Arinda A, Nakitende G, Otim L. Timing and reasons for coming late for the first antenatal care visit by pregnant women at Mulago hospital, Kampala Uganda. *BMC Pregnancy Childbirth*. 2013;13(1):121.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

